

Abstract: Determination of adatom interaction energies by a Monte Carlo calculation: Oxygen on W(110)^{a)}

E. D. Williams,^{b)} S. L. Cunningham, and W. H. Weinberg^{c)}

Division of Chemistry and Chemical Engineering, California Institute of Technology, Pasadena, California 91125

(Received 19 September 1977; accepted 22 November 1977)

PACS numbers: 68.40.+e

A great deal of experimental data on the order-disorder transition of the $p(2 \times 1)$ structure of oxygen on W(110) is available.¹⁻⁴ Previous attempts to determine the adatom-adatom interaction energies from different sets of these data have been made.^{4,5} However, in both attempts a model for the interaction energies which was inconsistent with the symmetry of the surface was used.

The locations of the oxygen atoms in the $p(2 \times 1)$ structure on W(110) as determined by dynamical calculations of LEED I - V curves are shown in Fig. 1. The parameters ϵ_x denote the interaction energies between oxygen atoms which are separated by the arrows as shown. In the previous studies^{4,5} the parameters ϵ_a , ϵ_b , and ϵ_{2b} , where ϵ_a and ϵ_{2b} are attractive interactions and ϵ_b is repulsive, were used. But since the (10) and (01) directions are equivalent, it is a violation of symmetry considerations to assign different values to ϵ_a and ϵ_b . Furthermore, if this were the true set of interaction energies only the domain shown in Fig. 1 would occur and not the domain in which the rows of adatoms are in the (01) direction. Both the $(\frac{1}{2}, 0)$ and $(0, \frac{1}{2})$ spots are observed experimentally indicating that both domains are present. Therefore, this model for the interaction energies cannot be correct.

The parameters used in this study, ϵ_1 , ϵ_2 , ϵ_3 , and ϵ_4 , where ϵ_1 and ϵ_4 are attractive and ϵ_2 and ϵ_3 are repulsive, conform




FIG. 1. Schematic of the $p(2 \times 1)$ overlayer of oxygen on W(110) as determined by LEED along with two possible models for the oxygen-oxygen interaction energies: (1) ϵ_1 , ϵ_2 , ϵ_3 , and ϵ_4 which are used in this work; (2) ϵ_a , ϵ_b , and ϵ_{2b} which were used in Refs. 4 and 5.

to the symmetry of the surface and allow the formation of perpendicular domains. While ϵ_2 and ϵ_3 are undoubtedly different, they are set equal in this study since the data are insufficient to determine four parameters uniquely.

A Monte Carlo calculation using a 30×30 lattice with periodic boundary conditions was used to simulate equilibrium configurations of the adatoms as a function of temperature for various sets of interaction energies. The LEED beam intensities for these configurations were calculated within the kinematic approximation and the intensity versus temperature (I - T) curves thus generated were compared with the experimental data⁴ to determine the best set of parameters. I - T curves were calculated for a large number of ratios of $\epsilon_1:\epsilon_2 = \epsilon_3:\epsilon_4$ at half coverage ($\theta = 0.5$). The parameters were then scaled to give the correct transition temperature ($T_c = 730$ K) at half coverage, and I - T curves were calculated for quarter coverage ($\theta = 0.25$). It was possible to eliminate many sets of parameters as they gave an incorrect transition temperature at quarter coverage.




FIG. 2. Intensity of the $(\frac{1}{2}, 0)$ LEED spot for W(110)- $p(2 \times 1)O$ as a function of temperature for two different coverages. The theoretical curve calculated using $\epsilon_1 = -2.1$ kcal/mol, $\epsilon_2 = \epsilon_3 = 1.7$ kcal/mol, and $\epsilon_4 = -0.7$ kcal/mol is compared with the experimental data from Ref. 4.

The most satisfactory ratio of parameters is $\epsilon_1:\epsilon_2:\epsilon_4 = -1:5/6:-1/3$. The scaling factor which gives $T_c = 730$ K is 2.09 kcal/mol so that the final values are:

$$\epsilon_1 = -2.1 \text{ kcal/mol}$$

$$\epsilon_2 = \epsilon_3 = 1.7 \text{ kcal/mol}$$

$$\epsilon_4 = -0.7 \text{ kcal/mol.}$$

Figure 2 shows a comparison of the I - T curves calculated using these parameters with the experimental data. The fit at $\theta = 0.53$ for temperatures below T_c is excellent. The poor fit above T_c is not considered important as the shape of the curve above the transition temperature is a strong function of lattice size.^{7,8}

The values of the interaction energies determined in this study are of the order of magnitude expected.⁹ They also show decrease in magnitude with distance, as opposed to earlier studies^{4,5} in which ϵ_{2b} , the interaction energy between adatoms two lattice spacings apart was equal to ϵ_a , the interaction energy over one lattice spacing. Most important, these pa-

rameters are consistent with the symmetry requirements of the system.

Acknowledgment. This work was supported in part by the AROD under Grant No. DAHCO4-75-0170.

^a)Complete paper to appear in J. Chem. Phys.

^b)National Science Foundation Predoctoral Fellow.

^c)Alfred P. Sloan Foundation Fellow, and Camille and Henry Dreyfus Foundation Teacher-Scholar.

¹)J. C. Tracy and J. M. Blakely, Surf. Sci. **15**, 257 (1969).

²)J. C. Buchholz and M. G. Lagally, Phys. Rev. Lett. **35**, 442 (1975).

³)T. Engel, H. Niehus, and E. Bauer, Surf. Sci. **52**, 237 (1975).

⁴)T.-M. Lu, G.-C. Wang, and M. G. Lagally, Phys. Rev. Lett. **39**, 411 (1977).

⁵)G. Ertl and D. Schillinger, J. Chem. Phys. **66**, 2569 (1977).

⁶)M. A. Van Hove and S. Y. Tong, Phys. Rev. Lett. **35**, 1092 (1975).

⁷)D. P. Landau, Phys. Rev. B **13**, 2997 (1976).

⁸)G. Doyen, G. Ertl, and M. Plancher, J. Chem. Phys. **62**, 2957 (1975).

⁹)T. L. Einstein and J. R. Schrieffer, Phys. Rev. B **7**, 3629 (1973).